

BUKTI KORESPONDENSI

Lampiran	:	Peer review proses korespondensi submit Publikasi Jurnal Ilmiah Nasional Terakreditasi
Nama Jurnal	:	Jurnal Farmasi Sains Vol. 7 No.1 2021 Hal: 74-80
Index	:	Terakreditasi SINTA 3
Judul Jurnal	:	Irritation Test of Bawang Dayak (<i>Eleutherine bulbosa</i> (Mill.) Urb.) Extract Cream with Human Patch Test Method

No	Item	Tanggal	Halaman
1	Register akun jurnal dan submit artikel	23 Maret 2021	1
2	Review Process	18 April 2021	2
3	Article for final proof (Copyediting)	28 April 2021	5
4	Article Published	01 Mei 2021	6
5	Ethical approval	09 April 2020	9

1. Register akun jurnal dan submit artikel

The screenshot displays the submission management interface for the journal 'Jurnal Farmasi Sains dan Praktis'. The user is logged in as 'syahrida'. The submission title is 'IRRITATION TEST OF BAWANG DAYAK (Eleutherine bulbosa(Mill.) Urb.) EXTRACT CREAM WITH HUMAN PATCH TEST METHOD' by Syahrida Dian Ardhany, Susi Novaryatiin, Mohammad Rizki Fadhil Pratama, and Zulkhurnain U... The submission is currently in the 'Production' stage, with other stages being 'Submission', 'Review', and 'Copyediting'. Under 'Submission Files', two files are listed: '14114-1 syahrida, New 2020 template JFSP.doc' and '14435-1 elmiawati, Syahrida_Send Reviewer 1 April.doc', both identified as 'Article Text'. A 'Download All Files' button is available. The 'Pre-Review Discussions' section is currently empty, showing 'No Items'.

Table 1. *Beringin* bark extract cream formulation

Materials	5% <i>Beringin</i> bark	20% <i>Beringin</i> bark	Cream base
<i>Beringin</i> bark ethanol extract	1250 mg	5000 mg	
Carboxymethyl cellulose	2000 mg	2000 mg	2000 mg
Oil phase			2000 mg
Stearic acid	5000 mg	5000 mg	5000 mg
Alkyl lactate	750 mg	750 mg	750 mg
Paraffin liquid	6250 mg	6250 mg	6250 mg
Aqueous phase			
Tetrasodium EDTA	375 mg	375 mg	375 mg
Triethylamine	25 mg	25 mg	25 mg
Water	30 mL	30 mL	30 mL
Total	20000 mg	20000 mg	20000 mg

Patch Test Study
A total of 20 volunteers were recruited for the patch test study, and each volunteer used two hands for the base and cream test. The first study for the cream test (right hand) consisted of 10 subjects, including nine normal (1) women with an age range of 19-23 years were living in Palembang Kota city, while the second study for the base test with the same volunteers (left hand). A patch test was done on the forearm of each volunteer on the first day of skin testing. The test was conducted by applying cream and base on the forearm with a diameter of 2 cm, 100 mg of the base, and 100 mg of cream formulation applied on the left and right forearm of the volunteers which was previously marked with a size of 4 x 2.5 cm, respectively. The method of application of cream to the volunteer forearm is shown in Figure 1. After applying base and cream formulation, it was then enclosed with the help of surgical paper bandages, as shown in Figure 2. Then the patch was left to stand for 24 hours to observe for any irritation on the volunteers' skin. The duration of contact for 24 hours was expected to be optimal in observing the irritation test of topical preparations on the skin (Hidayat et al., 2014). After 24 hours, the patch test was removed, and the forearm was washed with normal saline solution, after which erythema was determined by using a score scale (Akhbar et al., 2014) as presented in Table 2.

Table 2. Skin reaction scoring criteria

Score	Reaction
0	No response
1	Mild redness/erythema
2	Erythema with slight to moderate edema
3	Vesicles (small blisters) or papules (small, circumscribed elevations)
4	Bullae (large blisters), necrosis, or other severe reaction

3. RESULTS AND DISCUSSION
Beringin bark is one of the typical plants of Central Kalimantan, which has several secondary metabolites such as flavonoids, alkaloids, saponins, and tannins (Sugeng et al., 2014).

Ashlagi, 2019. Based on some literature, these metabolites have activities that can inhibit the bacteria that cause acne with their respective mechanisms, such as flavonoids with antimicrobial mechanisms that generally synergize with antibacterial activity (Prasetyo et al., 2012; Kim et al., 2014; Ghoshal et al., 2019; Mansour & Elghayour, 2020; Komar et al., 2020; Prasad et al., 2020). Other studies have also reported that other secondary metabolites from *Beringin* bark, such as ginsenosides, are known to inhibit cell wall synthesis from bacteria (Chang & Aziz, 2019). This is very important to know before making a preparation. The human patch test results showed that neither the formulations used nor the cream base showed an irritating reaction to all the volunteers involved. The visualization of the results of cream application to volunteer samples is presented in Figure 3 to 5, while a recapitulation of the skin reaction scores is presented in Table 3.

Table 3. The results of irritation tests on volunteers

Sample	Volunteer skin irritation scores																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
5% <i>Beringin</i> bark cream	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20% <i>Beringin</i> bark cream	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cream base	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

In previous studies, the primary irritation test for *Beringin* bark extract cream was carried out on the rabbit, showing that it does not irritate the test animal skin (Akhbar et al., 2019). This study is then continued to the evaluation stage of skin irritation in humans as one of the essential things a product can be used by the public, especially cosmetics for long-term use (Ali et al., 2017). The method used in this study is the human patch test. Patch testing helps confirm the presence of an allergy and identify the most common allergens and cosmetic products, causing dermatitis (Komar et al., 2020).



Figure 3. Appearance of irritation test results on the forearm (A) before and (B) after application of cream base for 24 hours

Figure 1. Application of cream on the forearm



Figure 2. The site of application of the cream is covered with a post-top bandage



Figure 3. Appearance of irritation test results on the forearm (A) before and (B) after application of 5% *Beringin* bark extract cream for 24 hours



Figure 4. Appearance of irritation test results on the forearm (A) before and (B) after application of 20% *Beringin* bark extract cream for 24 hours



Figure 5. Appearance of irritation test results on the forearm (A) before and (B) after application of cream base for 24 hours

Apart from visual observation of irritation, the volunteers were also interviewed verbally to determine a particular sensation in each treatment. The interview results will provide information about irritation symptoms that are not visible visually, including an itching and pain sensation (Gagang et al., 2017; Daud et al., 2018). From interviews with all volunteers, it was found that the two cream formulations did not cause itching or other sensations who report a slight itching from the use of the cream base. However, the itching sensation was reported to be minimal and did not cause the feeling of scratching the area. The itching sensation may be due to unfamiliar volunteers with a cream base that contains a fair amount of oil phase. Cream preparations contain a relatively large amount of oil phase, especially those with the water in oil (W/O) type (Makhrum et al., 2017). In contrast to other topical preparations such as gels, which contain more water and rarely cause an itching sensation (Pati et al., 2019). However, cream preparations will provide an advantage of longer contact time between the active ingredients and the skin (Gustawanto et al., 2017). Although it seems simple, the irritation test for cream preparations made from active plant extracts against humans by the patch test method is rarely reported. Some results only report the results of irritation tests on test animals like mice and rabbits, as reported by Ali et al. (2017) and Lakshmi et al. (2019). In humans, irritation tests of cream with active extracts of natural ingredients have previously been reported on *Ficus sp.* (Khan et al., 2019), *Asplenium bilobatum* (Suryaningrum et al., 2018), *Mangifera indica* (Shah & Siddiqui, 2017), and *Cassia toria* (Munawar et al., 2017), all of which reported no irritation to the skin of volunteers. Meanwhile, the irritation test on human volunteers from a cream made from *Beringin* bark has never been reported before. This study's results are expected to provide valuable information in the development of cream preparations made from *Beringin* bark extract, especially as an anti-acne cream.

4. CONCLUSION
The results of this study indicate that the *Beringin* bark extract cream with a concentration of 5 and 20% w/w does not irritate human skin. This study is a continuation of previous studies to get additional data to produce a suitable formulation without causing harmful effects on the skin. Further research is needed to develop testing effectiveness cream


of *Beringin* bark in humans with acne faces to produce an anti-acne product with reasonable safety and high quality.

5. ACKNOWLEDGMENT
This research was funded by an internal grant from Universitas Muhammadiyah Palembang.

6. CONFLICT OF INTEREST
The authors declare that there is no competing conflict of interest.

7. REFERENCES
Akbar N, Khan H.M.S., Ashraf S, Muhammad J.S., Ali A. (2014). Skin depigmentation activity of *Croton tiglium* extract cream. *Tropical Journal of Pharmaceutical Research*, 13(11):1803-1809.
Ali A., Akbar N., Munzir A.M., Khan M.S., Iqbal F.M., Zaidi S.S. (2013). In vivo skin irritation potential of a cream containing *Moringa oleifera* leaf extract. *Asian Journal of Pharmacy and Pharmacology*, 7(6):289-291.
Ashlagi S.D., Liliandari R.R., Supriatna S. (2019). Uji Efikasi Formulasi Sediaan Krim Ekstrak *Beringin* (*Glycyrrhiza lewneyi* HBK.) UMS pada Kulit. *Abstrak Patch Test*. *Jurnal Dermatologi*, 5(1):63-69.
Ashlagi S.D., Supriatna S. (2019). Antibacterial activity of ethanolic extract *Beringin* bark (*Glycyrrhiza lewneyi* HBK.) in cream against *Propionibacterium acnes*. *International Journal of Applied Pharmaceutics*, 11(Special Issue 9):3-4.
Cantaria T.M., Chen H., Elmaghrabi B.F. (2016). Optimizing Non-Antibiotic Treatments for Patients with Acne: A Review. *Dermatology Therapy*, 6(4):355-78.
Daud N.S., Akbar A.J., Suniliana P., Kamaliyah D. (2018). Formulation of Seed Slime (*Arachis sativa*) Anti-Acne Emulsion using Tween 80-Span 80 as Emulsifying and HPMSC as Gelling Agent. *Review Journal of Pharmacy*, 12(1):61-67.
Elmaghrabi B.F., Mohsen A.M.E., Ibrahim I.M., Mubayyidh M.H., Elmaghrabi M.E. (2016). Effectiveness of topical preparation oil on propionibacterium. *Journal of Pharmaceutical Clinical, Cosmetic and Investigational Dermatology*, 9:333-8.
Fung M.A., Sulandra H.I., Andriani K.S., Lakshmi R.M., Kati P., Ryan C., Ely J., Kanti A. (2011). Historical perspective on the use of visual grading scales in evaluating skin irritation and sensitization. *Contact Dermatitis*, 65(2):65-75.
Fritsch Anwar M.G.D., Elmaghrabi F., Yildirim W., Samir L.C.D., Bravetti I.L., Sotomayor C.E., Baggio T.M. (2012). Correlation among antimicrobial, antimutagenic, hemolytic, and antiproliferative properties of *Lepidium sibiricum* extract. *International Journal of Molecular Sciences*, 13(7):9260-77.
Göktaşlı S., Demirezen S., Kılıncıoğlu İ. (2019). Comprehensive review of antimicrobial activities of plant flavonoids. *Phytochemistry Review*, 18:241-72.
Hassan A., Usha B., Inam M. (2017). The antioxidant activity and phytochemical analysis of medicinal plant *Vernonia biloba*. *Letters in Applied Microbiology*, 64:732-738.

Hesta K., Tanjung C., Yacobina M., Ghani T., Mawanga K. (2014). Study of the usefulness of patch testing and use test to predict the safety of commercial topical drugs. *The Journal of Dermatology*, 41(6):605-613.
Khan H., Akbar N., Ali A. (2014). Effects of Cream Containing *Ficus sp.* Fruit Extract on Skin Parameters: In vivo Evaluation. *Indian Journal of Pharmaceutical Sciences*, 76(6):540-544.
Kim S.B., Noh H.B., B.S., Lee S.H., Kim J.M., Choi C.W., Bae S.P. (2018). In Vivo Antioxidant and Anti-Propionibacterium acnes Activities of Cold Water, Hot Water, and Medicated Extracts, and Their Respective Ethyl Acetate Fractions, from *Sargassum thunbergii* L. *Biomol Biotechnol*, 21(11):3001.
Komar P., Pratiwi R. (2014). Patch Testing in Suspected Allergic Contact Dermatitis to Cosmetics. *Dermatology Research and Practice*, 2014:69537.
Kumar K., Mishra R.C., Yadav R. (2020). Antimicrobial and cytotoxic studies of *Acacia gyaladica* root extract and their green emulsional silver nanoparticles. *Letters in Applied Microbiology*, 70(2):193-199.
Lakshmi R., Yezliya V., Ganesan S., Dines K., Zainab T., Zeynep N., Edaga G., Kanubhan A., Shobana V., Mahalingam A. (2019). Phytochemical analysis and evaluation of skin irritation, acute and sub-acute toxicity of *Cymbopogon citratus* essential oil in mice and rabbits. *Toxicology Reports*, 6:1298-1298.
Maha A. (2017). Acne vulgaris. *Medicine*, 65(6):386-389.
Mansour G., Elmaghrabi B., Gabr M., Eln S., Kamel A.C., Zayed F. (2019). Tannin profile, antioxidant properties, and antimicrobial activity of extracts from two Mediterranean species of pennis plant *Cymbopogon*. *BMC Complementary and Alternative Medicine*, 19:82.
Mansour Y., Elmaghrabi B. (2020). Chemical composition and antibacterial activity of methanolic extract from *Glycyrrhiza lewneyi* on typical foodborne pathogens. *Letters in Applied Microbiology*, 70(1):849-852.
Makhrum M., Liliandari R., Baghel C., Fatah S., Ternate I., Citra M. (2017). Formulation and evaluation of a water-in-oil cream containing herbal active ingredients and folic acid. *Drug Metabolism*, 9(2):121-129.
Nicoventra S., Ashlagi S.D. (2019). The Antibacterial Activity of *Beringin* Bark (*Glycyrrhiza lewneyi* HBK.) from Central Kalimantan Against Acne-Causing Bacteria. *International Journal of Applied Pharmaceutics*, 11(Special Issue 5):22-25.
Nicoventra S., Ashlagi S.D. (2020). Potential Anti-acne *Beringin* Bark (*Glycyrrhiza lewneyi* HBK.) LHA from Central Kalimantan Indonesia. *Pharmaceutical Journal*, 12(1):52-57.
Oon H.H., Wang S.N., Yee A.W.D.C., Cheong M.K., Goh C.L., Tan H.H. (2016). Acne Management Guidelines by the Dermatological Society of Singapore. *The Journal of Clinical and Aesthetic Dermatology*, 12(1):16-21.
Othman L., Hassan A., Abdel-Mansour R.M. (2019). Antimicrobial Activity of Polyphenols and Alkaloids in *Melastoma* Plants. *Phytochemistry in Microbiology*, 10:111.
Othman L.A., Othman L.U., Adnan A.G., Othman O.J., Othman A. (2019). Essential oil of *Albizia leonardus* Thunberg: phytochemical, antimicrobial and toxicity evaluations. *Letters in Applied Microbiology*, 67(1):61-63.
Prasad A., Krishna S.B.N., Seetha P., Pillay K., Ganesan R. (2010). Phytochemical composition and biological investigation of *Glycyrrhiza lewneyi* leaf. *Seed extract*. *Letters in Applied Microbiology*, 52(1):111-116.
Rizkani L., Dossoussi C. (2011). Recent advances in understanding *Propionibacterium acnes* (*Staphylococcus acnes*) in acne. *F1000Research*, 1:1000 Faculty Rev-193.



Respon terhadap Komentar Reviewer

Judul artikel : IRRITATION TEST OF BAWANG DAYAK (Eleutherine bulbosa (Mill.) Urb.) EXTRACT CREAM WITH HUMAN PATCH TEST METHOD

Reviewer I

Komentar Reviewer	Respon Author
1. Reason for using concentrations of 5% and 20%.	1. Previous studies have made creams with 5,10,15 and 20% concentration of bawang dayak and all concentrations of creams did not show irritation in the test animals. So that in this study it is enough to use the lowest (5%) and highest (20%) concentrations only
2. The research method design has not been described. For example: open label and pre post controlled design, etc.	2. Study design of this study is pre post-test controlled design (fixed in the manuscript)
3. Inclusion and exclusion criteria has not described	3. The inclusion criteria of this study are healthy person, 18-30 years old, there is no history of allergic-related illness, cooperate well and wiling to be volunteer for this study. The exclusion criteria of this study were an unhealthy person (excessive sweating, the skin has a wet wound, or unnormal skin). (fixed in the manuscript)
4. Were there any checks carried out to show that the subject was healthy?. Explain if an examination was carried out	4. No examination carried out only physically
5. The figure and title were made on one page	5. fixed in the manuscript

Komentar Editor (jika ada)

Komentar Reviewer	Respon Author
-	• -

Revisi lain yang dilakukan oleh penulis (jika ada)

-

3. Article for final proof (Copyediting)

The screenshot displays the submission management interface for the journal 'Jurnal Farmasi Sains dan Praktis'. The article title is 'IRRITATION TEST OF BAWANG DAYAK (Eleutherine bulbosa(Mill.) Urb.) EXTRACT CREAM WITH HUMAN PATCH TEST METHOD' by Syahrida Dian Ardhanry, Susi Novaryatlin, Mohammad Rizki Fadhil Pratama, and Zulkhurnain U... The interface is currently in the 'Copyediting' stage, with tabs for 'Submission', 'Review', 'Copyediting', and 'Production'. Below the article title, there are two 'Copyediting Discussions' from user 'elmiawati' on April 28 and April 29, both with 0 replies. At the bottom, a 'Copyedited' section shows a file named '14936-2 editor_psp_10_Syahrida 74-80.pdf (2)' with the label 'Article Text'.

4. Article Published

Jurnal Farmasi Sains dan Praktis
ABOUT ▾ CURRENT ARCHIVES ANNOUNCEMENTS

Q. SEARCH

HOME / ARCHIVES / VOL 7 NO 1 (2021) / ARTICLES

Check for updates

Section Articles

IRRITATION TEST OF BAWANG DAYAK (*Eleutherine bulbosa*(Mill.) Urb.) EXTRACT CREAM WITH HUMAN PATCH TEST METHOD

https://doi.org/10.31603/pharmacy.v7i1.4854

■ Syahrida Dian Ardhanay
 chass501@gmail.com (Primary Contact)
 Muhammadiyah University of Palangkaraya
<https://orcid.org/0000-0002-8606-8991>

■ Mohammad Rizki Fadhil Pratama
 Muhammadiyah University of Palangkaraya
<https://orcid.org/0000-0002-0727-4392>


■ Susi Novaryatiin
 Muhammadiyah University of Palangkaraya
<https://orcid.org/0000-0003-0696-6546>

■ Zulkhurnain Utar
 Malaysian Institute of Pharmaceuticals and Nutraceuticals (IPharm),
 National Institute of Biotechnology Malaysia (NIBM)



Abstract


Topical agents indicated for the treatment of acne have the potential to cause irritation or allergic contact dermatitis. This study investigates the irritancy potential of anti-acne cream of bawang dayak (*Eleutherine bulbosa* (Mill.) Urb.) previously tested for microbiological effectiveness with the lowest concentration of 5% and the highest concentration of 20%. The method used in this study is the human



JFSP Vol.7, No.1, April 2021, Hal. 74-80 pISSN: 2549-8068 eISSN: 2579-4558

Jurnal Farmasi Sains dan Praktis
(JFSP)

<http://journal.umng.ac.id/index.php/pharmacy>



IRRITATION TEST OF BAWANG DAYAK (*Eleutherine bulbosa* (Mill.) Urb.) EXTRACT CREAM WITH HUMAN PATCH TEST METHOD

Syahrida Dian Ardhanay^{1*}, Susi Novaryatiin¹, Mohammad Rizki Fadhil Pratama¹, Zulkhurnain Utar²

¹ Department of Pharmacy, Faculty of Health Sciences, Muhammadiyah University of Palangkaraya, Palangka Raya, 73111, Indonesia

² Malaysian Institute of Pharmaceuticals and Nutraceuticals (IPharm), National Institute of Biotechnology Malaysia (NIBM), Ministry of Science, Technology and Innovation (MOSTI), Putrajaya, 43700, Malaysia

Submitted: 23-03-2021
 Revised: 01-04-2021
 Accepted: 22-04-2021

Corresponding author:
 chass501@gmail.com

ABSTRACT
 Topical agents indicated for the treatment of acne have the potential to cause irritation or allergic contact dermatitis. This study investigates the irritancy potential of anti-acne cream of bawang dayak (*Eleutherine bulbosa* (Mill.) Urb.) previously tested for microbiological effectiveness with the lowest concentration of 5% and the highest concentration of 20%. The method used in this study is the human patch test. A total of 20 volunteers were recruited for the patch test study, using the cream. The results showed that all volunteers did not experience irritation both in the 5% or 20% bawang dayak extract cream formulations. However, the interview results were found that some volunteers experienced a slight itching without any significant skin adverse reactions on the cream application. Therefore, based on these initial findings it can be safely concluded that the cream of bawang dayak does not cause significant skin adverse reactions and good enough for further development for anti-acne cream dosage form.

Keywords: Acne, Bawang dayak, Cream, Patch test

1. INTRODUCTION

Acne vulgaris is one of the skin issues that can affect anyone. Acne vulgaris is a disease that causes inflammatory (papules, pustules, or nodules) or non-inflammatory lesions (open and closed comedones) (Tan et al., 2018). Typically located on the forehead, upper chest, or back with the highest density of sebaceous follicles (Malito, 2017). Acne vulgaris may affect all age groups but are typically experienced by adolescents who experience puberty when hormonal changes occur at 85% (Savovic et al., 2017). While acne is a common skin issue, sometimes it can also make adolescents feel uneasy and make a significant contribution to psychological distress. The research has shown that acne faced by adolescents tends to reduce adolescent self-esteem, making them insecure or anxious (Oon et al., 2019; Prabhakar et al., 2020).

Propionibacterium acnes are Gram-positive bacteria, the normal anaerobic flora of the skin that is an essential component of the skin microbiota (Platsidaki et al., 2018). Administering antibiotics for acne treatment is most frequently done to minimize the bacterial population. However, excessive use of antibiotics may lead to bacterial resistance. Increasing the use of antibiotics worldwide without the development of new antibiotics is a very urgent problem (Cisarova et al., 2016). Therefore, it is important to consider or turn from now on to using non-antibiotic preparations as far as possible (Rathi, 2011), such as exploring local anti-acne plants. Based on previous research, an ethanolic extract of bawang dayak and cream formulation have antibacterial activity against acne-causing (*P. acnes*, *Staphylococcus epidermidis*, and *Staphylococcus aureus*), the result of the minimum inhibitory concentration (MIC) test was 0.19% (Novaryatin & Ardhanay, 2019; Novaryatin & Ardhanay, 2020). The inhibition caused by

the ethanolic extract of bawang dayak which has been tested for phytochemical compounds containing secondary metabolites such as alkaloids, flavonoids, tannins, and saponins which have antioxidant and antibacterial potential activity (Gorniak et al., 2019; Othman et al., 2019; Maisetta et al., 2019; Tagousoop et al., 2018; Yazdani et al., 2019; Hassan et al., 2019). Besides, the primary irritation test results in rabbits showed the ethanolic extract of bawang dayak did not irritate topically (Ardhanay et al., 2019). Based on this background, it is necessary to research the irritation test on human skin further using the human patch test method to determine whether the cream is safe to use on human skin before it is used as anti-acne.

2. METHODS

Plant Collection and Identification

Fresh bulbs of bawang dayak were collected from the farmer in Sei Gohong Bukit Batu, Palangka Raya, Central Kalimantan and authenticated by Dr. Joeni Setijo Ralajoe from Indonesian Institute of Sciences Research Center for Biology, Bogor Indonesia.

Preparation of Plant Extract

The bawang dayak were cut into pieces and dried at a temperature of not more than 40°C. The dried bulb is then crushed with a grinder to become a powder. The powder of the bulbs bawang dayak was extracted with 96% ethanol using a percolator until the solvent's color returned colorless, and once the process was finished, all extracts were concentrated in a rotary evaporator.

Formulation of Cream Preparation

Bawang dayak bulbs after the extraction process were formulated as creams by different concentration, the lowest (5%) and the highest (20%) based on the study before (Ardhanay & Novaryatin, 2019) with adding cinnamon, honey, and aqua Menthae piperitae to improve the odor of cream formulation as shown in Table 1. Menthae piperitae also can give a better creamy smell, so it is more comfortable to use (Elsaie et al., 2016).

Table 1. Bawang dayak extract cream formulation

Materials	5% bawang dayak extract cream	20% bawang dayak extract cream	Cream base
Bawang dayak ethanolic extract	1250 mg	5000 mg	-
Cinnamon powder	2500 mg	2500 mg	2500 mg
Honey	2000 mg	2000 mg	2000 mg
Oil phase			
Stearic acid	5000 mg	5000 mg	5000 mg
Adeps lanae	750 mg	750 mg	750 mg
Paraffin liquid	6250 mg	6250 mg	6250 mg
Aqueous phase			
Triethanolamine	375 mg	375 mg	375 mg
Nipagin	25 mg	25 mg	25 mg
aqua Menthae piperitae	20 mL	20 mL	20 mL
Aquades ad	25000 mg	25000 mg	25000 mg

Study Design

This study was approved by the Health Research Ethics Committee of Universitas Aisyiyah Yogyakarta with ethical approval No. 1638KEP-UNISA/Y/2020. Study design of this study is pre-post-test-controlled design. Written informed consent was obtained from all participants. The inclusion criteria of this study were a healthy person, 18-30 years old, there is no history of allergic-related illness, cooperate well and willing to be volunteer for this study. The exclusion criteria of this study were an unhealthy person (excessive sweating, the skin has a wet wound, or unusual skin). Patch test of cream ethanolic extract Bawang dayak was

conducted on 20 healthy individuals used two hands each participant, one for control and one for the cream test. The 20 volunteers tested at the same time and concentration both base and cream test, each 10 volunteers for cream with 5% w/w or 20% w/w on right hand and base on left hand. The volunteers will be informed to don't excessive activities causing sweat.

Patch Test Study

A total of 20 volunteers were recruited for the patch test study, and each volunteer used two hands for the base and cream test. The first study for the cream test (right hand) consisted of 20 subjects, including nine men and 11 women with an age range of 19-25 years were living in Palangka Raya city, while the second study for the base test with the same volunteers (left hand). A patch test was done on the forearm of each volunteer on the first day of skin testing. The test was conducted by applying cream and base on the forearm with a diameter of ± 3 cm, 100 mg of the base, and 100 mg of cream formulation applied on the left and right forearm of the volunteers which was previously marked with a size of 4×2.5 cm, respectively. The method of application of cream to the volunteer forearm is shown in Figure 1. After applying base and cream formulation, it was then enclosed with the help of surgical post-op bandages. Then the patch was left to stand for 24 hours to observe for any irritation on the volunteers' skin. The duration of contact for 24 hours was reported to be optimal in observing the irritation test of topical preparations on the skin (Horita et al., 2014). After 24 hours, the patch test was removed, and the forearm was washed with normal saline solution, after which erythema was determined by using a score scale (Aktar et al., 2014) as presented in Table 2.

Table 2. Skin reaction scoring criteria

Score	Reaction
0	No response
1	Well-defined erythema
2	Erythema with slight to moderate edema
3	Vesicles (small blister) or papules (small, circumscribed elevations)
4	Bullous (large blister), spreading, or other severe reaction

3. RESULTS AND DISCUSSION

Bawang dayak is one of the typical plants of Central Kalimantan, which has several secondary metabolites such as flavonoids, alkaloids, saponins, and tannins (Novaryatin & Arduhany, 2019). Based on some literature, these metabolites have activities that can inhibit the bacteria that cause acne with their respective mechanisms, such as flavonoids with antioxidant mechanisms that generally synergize with antibacterial activity (Freitas et al., 2012; Kim et al., 2018; Onuchere et al., 2019; Massoud & Reihaneh, 2020; Kumari et al., 2020; Penual et al., 2020). Other studies have also reported that other secondary metabolites from bawang dayak, such as eugenol are known to inhibit cell wall synthesis from bacteria (Pratama & Aziz, 2019). This is very important to know before making a preparation. The human patch test results showed that neither the formulations used nor the cream base showed an irritating reaction to all the volunteers involved. The visualization of the results of cream application to volunteer samples is presented in Figure 2, Figure 3, Figure 4, while a recapitulation of the skin reaction scores is presented in Table 3.

In previous studies, the primary irritation test for bawang dayak extract cream was carried out on the rabbit, showing that it does not irritate the test animal skin (Arduhany et al., 2019). This study is then continued to the evaluation stage of skin irritation in humans as one of the essential things a product can be used by the public, especially cosmetics for long-term use (Ali et al., 2013). The method used in this study is the human patch test. Patch testing helps confirm the presence of an allergy and identify the most common allergen and cosmetic products, causing dermatitis (Kumar et al., 2014).




Figure 4. Appearance of irritation test results on the forearm: (a) before; and (b) after application of cream base for 24 hours

Apart from visual observation of irritation, the volunteers were also interviewed verbally to determine a particular sensation in each treatment. The interview results will provide information about irritation symptoms that are not visible visually, including an itching and pain sensation (Zuuren et al., 2017; Daud et al., 2018). From interviews with all volunteers, it was found that the two cream formulations did not cause itching—only some volunteers who report a slight itching from the use of the cream base. However, the itching sensation was reported to be minimal and did not cause the feeling of scratching the area. The itching sensation may be due to unfamiliar volunteers with a cream base that contains a fair amount of oil phase. Cream preparations contain a relatively large amount of oil phase, especially those with the water in oil (W/O) type (Moldovan et al., 2017). In contrast to other topical preparations such as gels, which contain more water and rarely cause an itching sensation (Putri et al., 2019). However, cream preparations will provide an advantage of longer contact time between the active ingredients and the skin (Purnamawati et al., 2017).

Although it seems simple, the irritation test for cream preparations made from active plant extracts against humans by the patch test method is rarely reported. Some studies only report the results of irritation tests on test animals like mice and rabbits, as reported by Ali et al. (2013) and Lukidul et al. (2019). In humans, irritation tests of creams with active extracts of natural ingredients have been reported on *Ficus carica* (Khan et al., 2014), *Averrhoa bilimbi* (Sutrisnigah et al., 2018), *Mangifera indica* (Satria & Salsana, 2017), and *Cucumis longia* (Wattanakrai et al., 2007), all of which reported no irritation to the skin of volunteers. Meanwhile, the irritation test on human volunteers from a cream made from bawang dayak has never been reported before. This study's results are expected to provide valuable information in the development of cream preparations made from bawang dayak extract, especially as an anti-acne cream.

4. CONCLUSION

The results of this study indicate that the bawang dayak extract cream with a concentration of 5 and 20% w/w does not irritate human skin. This study is a continuation of previous studies to get additional data to produce a suitable formulation without causing harmful effects on the skin. Further research is needed to develop testing effectiveness cream of bawang dayak in humans with acne faces to produce an anti-acne product with reasonable safety and high quality.

5. ACKNOWLEDGMENT

This research was funded by an internal grant from Universitas Muhammadiyah Palangkaraya.

6. CONFLICT OF INTEREST

The authors declare that there is no competing conflicts of interest.

7. REFERENCES

Aktar N, Kham H.M.S., Ashraf S., Mohammad L.S., Ali A. (2014). Skin depigmentation activity of *Crocos sativus* extract cream. *Tropical Journal of Pharmaceutical Research*, 13(11),1803-1808.

Table 3. The results of irritation tests on volunteers

Sample	5% bawang dayak extract cream	20% bawang dayak extract cream	Cream base
1	0	0	0
2	0	0	0
3	0	0	0
4	0	0	0
5	0	0	0
6	0	0	0
7	0	0	0
8	0	0	0
9	0	0	0
10	0	0	0
11	0	0	0
12	0	0	0
13	0	0	0
14	0	0	0
15	0	0	0
16	0	0	0
17	0	0	0
18	0	0	0
19	0	0	0
20	0	0	0




Figure 1. (a) Application of cream on the forearm; and (b) the site of application of the cream is covered with a post-op bandage



Figure 2. Appearance of irritation test results on the forearm: (a) before; and (b) after application of 5% bawang dayak extract cream for 24 hours



Figure 3. Appearance of irritation test results on the forearm: (a) before; and (b) after application of 20% bawang dayak extract cream for 24 hours

- Ali A., Aktar N., Mumtaz A.M., Khan M.S., Iqbal F.M., Zaidi S.S. (2013). In vivo skin irritation potential of a cream containing *Moringa oleifera* leaf extract. *African Journal of Pharmacy and Pharmacology*, 7(6),289-293.
- Arduhany S.D., Effendi R.R., Novaryatin S. (2019). Uji Intra Formulasi Sediaan Krim Ekstrak Bawang Dayak (*Eleocharis bulbosa* (Mill.) Urb) pada Kelinci Albino Putih. *Jurnal Surya Medika*, 5(1),63-69.
- Arduhany S.D., Novaryatin S. (2019). Antibacterial activity of ethanolic extract Bawang dayak (*Eleocharis bulbosa* (Mill.) Urb) in cream against *Propionibacterium* species. *International Journal of Applied Pharmaceutics*, 11(Special Issue 5),1-4.
- Cinamon T.N., Chen E., Elewski B.E. (2016). Optimizing Non-Antibiotic Treatments for Patients with Acne: A Review. *Dermatologic Therapy*, 6(4),555-78.
- Daud N.S., Aldar A.J., Nuzuliana E., Kamalika (2018). Formulation of Saal Slices (*Achatina Falcata*) Anti-Acne Emulgel using Tween 80-Span 80 as Emulsifier and HPMC as Gelling Agent. *Borneo Journal of Pharmacy*, 1(2),64-67.
- Elsie L.T., Madsen A.M.E., Thrilum I.M., Moleky-Edlin M.H., Elsie M.L. (2016). Effectiveness of topical peppermint oil on symptomatic treatment of chronic pruritus. *Clinical, Cosmetic and Investigational Dermatology*, 9,333-8.
- Fargah M.A., Malhotra H.I., Andersen K.E., Lachapelle J.M., Kern P., Rymn C., Ely J., Kusti A. (2011). Historical perspective on the use of visual grading scales in evaluating skin irritation and sensitization. *Contact Dermatitis*, 65(2),65-75.
- Freitas Araújo M.G.D., Hilario F., Vilgas W., Santos L.C.D., Brunetti I.L., Sotomayor C.E., Banab T.M. (2012). Correlation among antioxidant, antimicrobial, hemolytic, and mutagenic properties of *Leitochia spiralis* leaves extract. *International Journal of Molecular Science*, 13(7),9260-77.
- Górnika I., Bartoszewski R., Królczewski J. (2019). Comprehensive review of antimicrobial activities of plant flavonoids. *Phytochemistry Review*, 18,241-72.
- Hasan A., Utih H., Isar M. (2019). The antioxidant activity and phytochemical analysis of medicinal plant *Veronica biloba*. *Letters in Applied NanoBioScience*, 8(4),732-738.
- Horita K., Tamone C., Yasoshima M., Ohtani T., Matsunaga K. (2014). Study of the usefulness of patch testing and use test to predict the safety of commercial topical drugs. *The Journal of Dermatol*, 41(6),505-513.
- Khan H., Aktar N., Ali A. (2014). Effects of Cream Containing *Ficus carica* L. Fruit Extract on Skin Parameters: In vivo Evaluation. *Indian Journal of Pharmaceutical Science*, 76(6),560-564.
- Kim S., Oh S., Noh H.B., Ji S., Lee S.H., Koo J.M., Choi C.W., Jhon H.P. (2018). In Vitro Antioxidant and Anti-Propionibacterium species Activities of Cold Water, Hot Water, and Methanol Extracts, and Their Respective Ethyl Acetate Fractions, from *Sanguisorba officinalis* L. Roots. *Molecules*, 23(11),3001.
- Kumar F., Paulose R. (2014). Patch Testing in Suspected Allergic Contact Dermatitis to Cosmetics. *Dermatology Research and Practice*, 2014,693387.
- Kumari R., Mishra R.C., Yadav J.P. (2020). Antioxidant and cytotoxic studies of *Acacia nilotica* twig extract and their green synthesized silver nanoparticles. *Letters in Applied NanoBioScience*, 9(2),975-985.
- Lukidul E., Tesfaye S., Gebreabdirou S., Dires K., Zembe T., Zegaye N., Feleke G., Kassahun A., Shiferaw Y., Mekonen A. (2019). Phytochemical analysis and evaluation of skin irritation, acute and sub-acute toxicity of *Cymbopogon citratus* essential oil in mice and rabbits. *Toxicology Reports*, 6,1289-1294.
- Mahro A. (2017). Acne vulgaris. *Medicine*, 45(6),386-389.
- Matsieta G., Batoni G., Caboni P., Eain S., Rinaldi A.C., Zucca P. (2019). Tannin profile, antioxidant properties, and antimicrobial activity of extracts from two Mediterranean species of parasitic plant *Cytisus*. *DMC Complementary and Alternative Medicine*, 19,82.
- Massoud Y., Reihaneh B. (2020). Chemical composition and antibacterial activity of methanolic extract from *Echinops robustus* on typical food-borne pathogens. *Letters in Applied NanoBioScience*, 9(1),849-852.
- Moldovan M., Lohman A., Bogdan C., Parasca S., Tomuta I., Crisan M. (2017). Formulation and evaluation of a water-in-oil cream containing herbal active ingredients and ferulic acid. *Clyd Medical*, 90(2),212-219.
- Novaryatin S., Arduhany S.D. (2019). The Antibacterial Activity of Bawang Dayak (*Eleocharis bulbosa* (Mill.) Urb) from Central Kalimantan Against Acne-Causing Bacteria. *International Journal of Applied Pharmaceutics*, 11(Special Issue 5),22-25.
- Novaryatin S., Arduhany S.D. (2020). Potential Anti-acne: Bawang Dayak (*Eleocharis bulbosa* (Mill.) Urb) from Central Kalimantan-Indonesia. *Pharmacognosy Journal*, 12(1),43-47.
- On H.H., Wong S.N., Wee Aw D.C., Cheong W.K., Goh C.L., Tam H.H. (2019). Acne Management Guidelines by the Dermatological Society of Singapore. *The Journal of Clinical and Aesthetic Dermatology*, 12(7),34-50.

- Othman L., Sleiman A., Abdel-Massih R.M. (2019). Antimicrobial Activity of Polyphenols and Alkaloids in Middle Eastern Plants. *Frontiers in Microbiology*, 10,911.
- Otusechere C.A., Durugbo E.U., Adesanya O., Omotolani O.F., Osbo A. (2019). Essential oil of *Alchornea laxiflora* (benfu): phytochemical, antimicrobial and toxicity evaluations. *Letters in Applied NanoBioScience*, 8(3),661-665.
- Perumal A., Krishna S.B.N., Sershen, Pillay K., Govender P. (2020). Phytochemical composition and biological investigation of *Trichilia emetica* Vahl. Seed extracts *Letters in Applied NanoBioScience*, 9(2),1111-1116.
- Platsidaki E., Desimioti C. (2018). Recent advances in understanding *Propionibacterium acnes* (*Cutibacterium acnes*) in acne. *F1000Research*, 7, F1000 Faculty Rev-1953.
- Prabhakar P.K., Nath D., Singh S., Mittal A., Baghel D.S. (2020). Formulation and evaluation of polyherbal anti-acne combination by using in-vitro model. *Biointerface Research in Applied Chemistry*, 10(1),4747-4751.
- Pratama M.R., Aziz I.R. (2019). Molecular Docking of Bawang Dayak (*Eleutheria bulbosa*) Secondary Metabolites as Bacterial Cell Wall Synthesis Inhibitor. In: Proceedings of the 1st International Conference on Science and Technology, Makassar, Indonesia. Ghent: EAI.
- Purnamaswati S., Indrawati N., Danarti R., Saeudin T. (2017). The Role of Moisturizers in Addressing Various Kinds of Dermatitis: A Review. *Clinical Medicine and Research*, 15(3-4),75-87.
- Putri A.N., Laila N., Forestyana D. (2019). Formula Optimization of *Ammonia muricata* Folium Ethanolic Extract of Anti Acne Gel Formulation using Factorial Design Method. *Borneo Journal of Pharmacy*, 2(2),63-70.
- Rathil S.K. (2011). Acne vulgaris treatment: the current scenario. *Indian Journal of Dermatology*, 56(1),7-13.
- Satria D., Siahman M.A. (2017). Formulasi Krim Anti - Aging dari Buah Mangga Mansalagi. *Jurnal Farmanesia*, 4(1),12-30
- Saviuc C., Ciubucă B., Dinică G., Bleoța C., Drumeș V., Chifiruc M.C., Popa M., Pircalabioru G.G., Marutescu L., Lazăr V. (2017). Development and Sequential Analysis of a New Multi-Agent, Anti-Acne Formulation Based on Plant-Derived Antimicrobial and Anti-Inflammatory Compounds. *International Journal of Molecular Science*, 18(1),175.
- Sotriingsih, Segal Z., Marhamah. (2018). Formulasi dan Uji Iritasi Gel Antibakteri dari Ekstrak Etanol Daun Belimbing Wuluh (*Averrhoa bilimbi* Linn) terhadap Bakteri *Staphylococcus aureus* dan *Pseudomonas aeruginosa*. *FAST Jurnal Sains Teknologi*, 2(1),1-9
- Tagousof C.N., Tamokou J.D., Kenyue I.C., Ngnokam D., Voutquenne-Nazabadioko L. (2018). Antimicrobial activities of saponins from *Melanthera elliptica* and their synergistic effects with antibiotics against pathogenic phenotypes. *Chemistry Central Journal*, 12,97.
- Tan A.U., Schlosser B.J., Paller A.S. (2018). A review of diagnosis and treatment of acne in adult female patients. *International Journal of Women's Dermatology*, 4(2),56-71.
- Wattanakrai P., Suvansachote S., Kulkollakam S., Rajatanavin N. (2007). The study of human skin irritation of a novel herbal skin care product and ingredients by human single closed patch testing. *Journal of the Medical Association of Thailand*, 90(6),1116-1122.
- Yazdani E., Talebi M., Zarsheenas M.M., Moein M. (2019). Evaluation of possible antioxidant activities of barberry solid formulation, a selected formulation from Traditional Persian Medicine (TPM) via various procedures. *Biointerface Research in Applied Chemistry*, 9(6),4517-4521.
- Zuuren E.J.V., Fedorowicz Z., Christensen R., Lavrijsen A., Arents B.W.M. (2017). Emollients and moisturisers for eczema. *Cochrane Database Systematic Reviews*, 2(2), CD012119.



UNIVERSITAS SARI MULIA
LEMBAGA PENELITIAN DAN PENGABDIAN KEPADA MASYARAKAT
DEWAN KOMITE ETIK PENELITIAN

Jl. Pramuka No.02 Banjarmasin Tlp. (0511) 3268105

Banjarmasin, 09 April 2020

No. SK : 031/KE-LPPM/UNISM/IV/2020
Lampiran : -
Perihal : Rekomendasi Penelitian

Peneliti yang disebutkan dibawah ini :

Ketua Peneliti : Syahrida Dian Ardhany
NIP/NIK/NIM : 14.0601.033
Anggota Peneliti : 1. Susi Novaryatiin
2. Mohammad Rizki Fadhil Pratama
3. Zulkhurnain Utar
Judul Penelitian : Uji iritasi kulit krim anti acne ekstrak Bawang Dayak
(*Eleutherine bulbosa* (Mill.) Urb.) metode *Human Patch Test*

Berdasarkan pertimbangan Dewan Komite Etik Penelitian diputuskan bahwa Peneliti yang disebutkan diatas telah **DISETUJUI** untuk melanjutkan penelitiannya.

Demikian surat persetujuan ini diterbitkan untuk dipergunakan dengan penuh tanggung jawab.

Menyetujui,

An. Ketua
Sekretaris Dewan Komite Etik Penelitian



H. Ali Rakhman Hakim, M.Farm., Apt
NIK. 1166012015073