



About us

Editorial board

Topics

Editorial policy / publishing ethics

Subscription information

Instruction to authors

Volumes

2022

The **INTERNATIONAL JOURNAL BIOAUTOMATION** is a continuation of the periodical Bioautomation, which was discontinued in the early nineties of the last century. It is an international journal for presentation of scientific results from the application of contemporary engineering and mathematical approaches and computer-oriented methods for the purposes of analysis, modelling, optimization and control.

2021

INT. J. BIOAUTOMATION comprises results in the main fields of:

- Bioinformatics
- Bioprocess systems
- Biomedical systems
- Biomedical physics
- Bioecological systems

2020

2019

2018

INT. J. BIOAUTOMATION accepts for publication original research papers, review articles, application notes, short communications and letters which give preliminary announcements of new theoretical and applied concepts. New experimental data, innovative experimental techniques or novel concepts in data correlation are also welcome. The journal is open to presentations from well known scientists, as well as to young researchers.

2017

2016

INT. J. BIOAUTOMATION is an open access journal offering everyone free and unlimited access to the full-text of peer-reviewed articles.

2015

The open access publishing policy contributes to publications' high visibility and easier fostering of new scientific collaborations between scientists from all over the world.

2014

All recommendations and remarks will be kindly accepted.

2013

Editorial Office

Institute of Biophysics and Biomedical Engineering
Bulgarian Academy of Sciences
105 Acad. Georgi Bonchev Str., room 211
1113 Sofia, BULGARIA

2012

2011

e-mail: bioautomation@biomed.bas.bg

2010

ISSN: 1314-2321 (on-line) 1314-1902 (print)

2009

© 2004-2022, BAS, Institute of Biophysics and Biomedical Engineering

2008

2007

2006

2005

2004

Shrink/Expand
all volumes

Indexed in:



SJR (2021): 0.2





[About us](#)
 [Editorial board](#)
 [Topics](#)
 [Editorial policy / publishing ethics](#)
 [Subscription information](#)
 [Instruction to authors](#)

Volumes

Editorial board

Honorary Chief Editor Stoyan Tzonkov, Bulgaria [+/- more]
 Chief Editor Mikhail Matveev, Bulgaria [+/- more]
 Deputy Chief Editor Tania Pencheva, Bulgaria [+/- more]

Topic Editors

Bioinformatics Krassimir Atanassov, Bulgaria [+/- more]
 Bioprocess systems Olympia Roeva, Bulgaria [+/- more]
 Biomedical systems Ivan Dotsinsky, Bulgaria [+/- more]
 Biomedical physics Iana Tsoneva, Bulgaria [+/- more]
 Bioecological systems Mitko Petrov, Bulgaria [+/- more]

Members

Alexander Petrov, Bulgaria [+/- more]
 Andon Kossev, Bulgaria [+/- more]
 Andrew Worth, Italy [+/- more]
 Antony Shannon, Australia [+/- more]
 Dawid Lochynski, Poland [+/- more]
 Denis Dochain, Belgium [+/- more]
 Enrico Spugnini, Italy [+/- more]
 Feng-Sheng Wang, Taiwan [+/- more]
 Giovanni Bortolan, Italy [+/- more]
 Govind Rao, USA [+/- more]
 Ilza Pajeva, Bulgaria [+/- more]
 Ivan Popchev, Bulgaria [+/- more]
 Ivaylo Christov, Bulgaria [+/- more]
 Ivo Iliev, Bulgaria [+/- more]
 Jean-Pierre Babary, France [+/- more]
 Jurgis Staniskis, Lithuania [+/- more]
 Maria Miteva, France [+/- more]
 Mihai Radu, Romania [+/- more]
 Nestor Vicente Torres, Spain [+/- more]
 Nicolai Christov, France [+/- more]
 Pratap Patnaik, India [+/- more]
 Rein Luus, Canada [+/- more]
 Rositsa Raikova, Bulgaria [+/- more]
 Stefan Hadjitodorov, Bulgaria [+/- more]
 Yordan Kostov, USA [+/- more]

Web Technology Petko Alov, Bulgaria [+/- more]

Web Management Elena Chorukova, Bulgaria [+/- more]
 Peter Vassilev, Bulgaria [+/- more]
 Olympia Roeva, Bulgaria [+/- more]

Shrink/Expand all volumes

Indexed in:



SJR (2021): 0.2





About us

Editorial board

Topics

Editorial policy / publishing ethics

Subscription information

Instruction to authors

Volumes

Volume 26, Issue 2

Indexed in:

2022

Identification of Metabolite Compounds and Biological Activity of *Diplazium esculentum*

131-140



SJR (2021): 0.2

- Volume 26, Issue 2
- Volume 26, Issue 1

Fathul Zannah, Mohamad Amin, Hadi Suwono, Betty Lukiati

doi: [10.7546/ijba.2022.26.2.000740](https://doi.org/10.7546/ijba.2022.26.2.000740)

[+/- how to cite] [+/- abstract] [full text]



2021

Validation of a Light Source for Phototoxicity in *in vitro* Conditions

141-152



Ivan Iliev, Inna Sulikovska, Elena Ivanova, Mashenka Dimitrova, Biliana Nikolova, Christina Andreeva

doi: [10.7546/ijba.2022.26.2.000837](https://doi.org/10.7546/ijba.2022.26.2.000837)

[+/- how to cite] [+/- abstract] [full text]

2020

2019

Evaluation of the Level of Alertness with Variations in Reaction Time in Bulgarian Sport Students

153-160



Mariya Zaharinova

doi: [10.7546/ijba.2022.26.2.000859](https://doi.org/10.7546/ijba.2022.26.2.000859)

[+/- how to cite] [+/- abstract] [full text]

2018

2017

Experimental Testing of a Prototype of an Active Elbow Orthosis Based on *in vivo* Investigation of Elbow Flexion/Extension of Healthy Subjects

161-174



Silvija Angelova, Emil Petrov, Plamen Raykov, Rositsa Raikova

doi: [10.7546/ijba.2022.26.2.000865](https://doi.org/10.7546/ijba.2022.26.2.000865)

[+/- how to cite] [+/- abstract] [full text]

2016

2015

A Clustering Algorithm for Tumor Gene Data Based on Improved DPC Algorithm

175-192



Wei Wang, Bo Gao

doi: [10.7546/ijba.2022.26.2.000872](https://doi.org/10.7546/ijba.2022.26.2.000872)

[+/- how to cite] [+/- abstract] [full text]

2014

2013

2012

A Key Protein Recognition Algorithm Based on High-order Complex Protein Network

193-208

Xiaojing Wang

doi: [10.7546/ijba.2022.26.2.000873](https://doi.org/10.7546/ijba.2022.26.2.000873)

[+/- how to cite] [+/- abstract] [full text]

2011

2010

2009

Sponsored by National Science Fund of Bulgaria, Grant No KP-06-NP3-37, 2022

2008

© 2022, BAS, Institute of Biophysics and Biomedical Engineering

2007

2006

2005

2004

Shrink/Expand all volumes

Identification of Metabolite Compounds and Biological Activity of *Diplazium esculentum*

Fathul Zannah^{1*}, Mohamad Amin², Hadi Suwono², Betty Lukiati²

¹University of Muhammadiyah Palangkaraya
Palangkaraya, Central Kalimantan, 63111, Indonesia
E-mail: fathulzannah.umpalangkaraya@gmail.com

²Department of Biology Education
State University of Malang
Malang, East Java, 65145, Indonesia
E-mails: mohamad.amin.fmipa@um.ac.id, hadi.suwono.fmipa@um.ac.id,
betty.lukiati.fmipa@um.ac.id

*Corresponding author

Received: November 12, 2020

Accepted: October 11, 2021

Published: June 30, 2022

Abstract: *Diplazium esculentum* is one of the medicinal plants used by Dayak tribes in Central Kalimantan to cure acne. The potential of *D. esculentum* can be proved through information on the active compounds of the extract obtained in decoction and infusa methods. Thus, a liquid chromatography-mass spectrophotometry (LC-MS)-based method is developed to identify the active compounds of *D. esculentum* extract in either decoction method or infusa method. The chemical compound potential of *D. esculentum* extract was then analysed using bioinformatics approach based upon the database of PASS online server. Results showed that the *D. esculentum* extract contained 81 chemical compounds in decoction method and 68 compounds in infusa method, in which the dominant compound was flavonoid. Moreover, PASS online web server analysis found 7 flavonoid compound groups potential as anti-acne containing antisebor, AR expression inhibitor and CYP1A1 inhibitor. This information could be very useful for designing a clinical test on plant natural compound potential for traditional drug development.

Keywords: Medicinal plant, Fern, Anti-acne, Decoction and Infusa methods.

Introduction

Vegetable fern *Diplazium esculentum* is a plant wildy growing in the river bank and highly humid garden. This plant is one of the medicinal plants used by Dayak tribe in Central Kalimantan to traditionally cure acne [35]. The ferns used as medicine can become source of good natural bioactive product that is potentially developed as new drug [4].

There are 4 factors affecting the occurrence of acne, epidermic follicular hyper-proliferation, increased sebum production, inflammation, and growth of *Propionibacterium acnes* production and follicular hyperkeratosis gives favourable pilobaceous condition for *P. acnes* growth [5].

Phytochemical screening of *D. esculentum* indicates the presence of polyphenol, alkaloid, tannin and saponin through ethanol extraction and flavonoid, polyphenol, alkaloid, tannin, and saponin in water extraction [36]. Flavonoid is only found in water extraction, decoction method and infusa method, but few information is available for active compounds of *D. esculentum* in water extract of both methods. To obtain information on chemical compounds of *D. esculentum* water extract, liquid chromatography-mass spectrometry

(LC-MS) is used, since this analysis can yield large spectral data [25]. This technique possesses specificity and very high sensitivity.

The use of water extraction is environmental friendly because the plant is not victimized and the organic solvent is substituted with easy, economic, sound, and safe alternative [31] compared with other chemical solvent, such as ethanol. Solvent determination and temperature can influence the extract quantity and quality of plant and its bioactive compounds [14].

Previous studies found the presence of relationship between flavonoid structure and its antibacterial activity [6]. Kaempferol as derivative of flavonoid in *Zingiber spectabile* Griff has antioxidant activity [26]. Luteolin, quercetin, and resveratrol also show antibacterial activity against *Staphylococcus aureus* [27]. Therefore, the potential of a medicinal plant is influenced by its chemical compounds and chemical structures.

This study aims to identify the active compounds of *D. esculentum* as herbal drug for acne treatment using LC-MS and determine the biological activity. The potential of *D. esculentum* as herbal medicine is recently unclear and has become major challenge to scientific approach, through *in vitro* and *in vivo* studies.

Materials and methods

Diplazium esculentum was collected from Palangkaraya, Central Kalimantan, Indonesia. Water was employed as solvent. Samples were cleansed and dried for about 5 days, finely ground, and extracted in distilled water using decoction method and infusa method. The former was carried out by putting one-hundred g of *D. esculentum* in 1000 mL of aquadest and boiled up to 90 °C, then filtered. The latter was done by pouring one-thousand ml of aquadest into a pan and boiled up to 90 °C, turned off the fire and added with 100 g of *D. esculentum* for 30 minutes, then filtered.

LC-MS analysis

Metabolite compound identification used Shimatsu LC-MS-8040 LC/MS. One μL of sample was injected into Shim Pack FC-ODS (2 mm D \times 150 mm, 3 μL) column with a capillary voltage of 3.0 kV and column temperature of 35 °C. The LC-MS analysis employed UPLC-MS equipped with binary pump. LC is connected to QTOF mass spectrometer joined with ESI. MS utilized positive ionization mode. The ESI parameter was set at the source temperature of 100 °C, 0.6 seconds scanning, and 80 minutes run time.

Biological activity analysis

A biological activity study is a basis required to set or select new molecule. The compound biological activity analysis utilized PASS online server [20]. It analyses the drug, biological activity, and various chemical compound targeting interactions. This server provides tools to select the compound of determined biological activity profiles including the pharmacotherapeutic effect desired and biochemical mechanism predicted as input information and provides electronic report on chemical compound literature. Biological potential analysis of compounds was carried out by inputting the chemical structures of *D. esculentum* extract obtained from LC-MS analysis in MOL file format or SDF file format on PASS online server. The outcome will show the biological potential of the compound based upon the database and accomplished with probability activity (PA) value, in which PA value ≥ 0.7 indicates that the computing prediction is not much different from the clinical test, while PA value ≤ 0.3 reflects that there is 50% probability similarity of the computing prediction to the clinical test. In the present study,

PASS online server analysis was directed to the compound potential of *D. esculentum* extract as antiseborrheic, AR expression inhibitor, and CYP1A1 inhibitor with PA \geq 0.7.

Results and discussion

This study found that the water extraction of *D. esculentum* contains various compounds as presented in Table 1.

Table 1. Number of metabolite compounds in water extraction of *D. esculentum* (decoction and infusa methods)

Method	Number of metabolite compounds
Decoction	81 compounds
Infusa	68 compounds

Table 1 demonstrates that the water extraction of *D. esculentum* in decoction method and infusa method contains various metabolite compounds – flavonoid, saponin, terpenoid, and carboxylic acid groups. *D. esculentum* extract holds more metabolite compounds in decoction method, 81 compounds, than those in infusa method, 68 compounds. This study found that water solvent is effective to use as organic solvent since it can filter various active compounds of *D. esculentum* extract, and difference in number of compounds indicates that decoction method is more effective than infusa method in water extraction. It could result from that compound solubility will rise with increasing temperature [3], boiling duration needs to be considered since increased boiling time in decoction method could result in decline in number of compounds in the extract [9].

The present study showed that dominant active compound of *D. esculentum* was flavonoid. Polyphenolic compounds produced by the plant and sent to human body through food and its occurrence in the tissue is dependent upon the intake of plant product [29]. Flavonoid has ability to induce human protecting enzyme system [17], such as anti-oxidant, anti-inflammation, anticancer, and cardiovascular protection [32]. Flavonoid extracted from *Cystoseira compressa* holds antibacterial activity as well [1], depending upon the structure, the substitution on the aromatic ring [33].

Water extraction of *D. esculentum* yields 31 flavonoid compound groups in decoction method (Table 2) and 24 compounds in infusa method (Table 3). There are more compounds of flavonoid group in decoction method than that in infusa method, since decoction method supports extraction better than infusa method and more appropriate for heat-resistant compounds, hard plant material, such as root, and will yield more soluble in oil than that in infusa method [2].

Table 2 shows that decoction method-based water extraction of *D. esculentum* contains 31 flavonoid compounds with the highest composition of calycosin 7 O β D glucoside and naringenin, 2.14473% and 2.01286%, respectively, and the lowest composition is procyanidin, 0.25975%. Naringenin has potential as antibacterial agent, especially against *Staphylococcus aureus* [19].

Table 3 demonstrates that infusa method-based water extraction of *D. esculentum* contains 24 flavonoid compounds with the highest composition of calycosin 7 O β D glucoside and naringenin, 2.134107% and 2.002882%, respectively, the lowest composition is recorded in

6 α hydroxymedicarpin, 0.423520%. More dominant flavonoid compound in water extraction of *D. esculentum* of both methods could result from high solubility of flavonoid in water, especially flavonoid glycoside [8].

Table 2. Metabolite compounds of flavonoid group in decoction method-based water extraction of *D. esculentum* (31 flavonoid compounds)

No	Compound	Molecular weight	Composition, %
1	Daidzein	254.24	0.40116
2	Coumestrol	268.22	0.95406
3	Biochanin B	268.26	1.62682
4	Apigenin	270.24	1.13494
5	Genistein	270.24	0.46703
6	Naringenin	272.25	2.01286
7	9 O methylcoumestrol	282.25	0.96241
8	Biochanin A	284.26	1.62318
9	Calycosin	284.26	1.13049
10	Homopterocarpin	284.31	1.02081
11	Luteolin	286.24	1.62337
12	6 α hydroxymedicarpin	286.28	0.42561
13	Quercetin	302.24	1.62634
14	Kaempferol 3 sulphate	365.29	1.10372
15	Daidzin	416.38	0.92221
16	Genistin	432.38	1.08384
17	Kaempferol 3 rhamnoside	434.35	0.98005
18	Calycosin 7 O β D glucoside	446.40	2.14473
19	Kaempferol 3 glucoside	448.38	1.49673
20	Kaempferol 7 O β D glucopyranoside	448.38	0.86479
21	Luteolin 7 glucoside	448.38	1.38711
22	Isoquercitrin	464.38	1.56773
23	Hyperoside	464.38	1.54309
24	Querciturone	478.36	0.85962
25	Leucocyanidin	306.27	0.78915
26	Luteolinidin 5 glucoside	432.38	0.37113
27	6'' O malonylgenistein	518.42	0.27540
28	Biochanin A 7 O β D glucoside 6'' O malonate	532.45	1.65780
29	Naringin	580.53	1.71495
30	Procyanidin	594.52	0.25975
31	Rutin	610.52	1.38667

Metabolite compounds of *D. esculentum* extract using either decoction method or infusa method could become very useful information for scientific prove in relation with the potentials as medicine. *D. esculentum* is one of the medicinal plants used by Dayak tribe in Central Kalimantan to traditionally cure acne [36]. It is important for traditional drug development from plant extract through ethnomedicinal usage history [15, 28].

Based on the results of previous studies, *D. esculentum* has several potentials. *D. esculentum*, a commonly consumed seasonal vegetable, has been reported to have some pathological

effects, especially on the male reproductive function. The study result show that *D. esculentum* has a potential as an antifertility agent after trial of male Swiss albino mouse [24]. Besides that, *D. esculentum* also potential as anti-coagulant agent [21] and as antioxidant and antidiabetic agent [13].

Table 3. Metabolite compounds of flavonoid in infusa method-based water extraction of *D. esculentum* (24 flavonoid compounds)

No	Compound	Molecule weight	Composition, %
1	Daidzein	254.24	0.658690
2	Coumestrol	268.22	0.949400
3	Biochanin B	268.26	1.618841
4	Apigenin	270.24	1.129313
5	Genistein	270.24	0.464718
6	Naringenin	272.25	2.002882
7	9 O methylcoumestrol	282.25	0.957634
8	Biochanin A	284.26	1.615133
9	Calycosin	284.26	1.124871
10	Homopterocarpin	284.31	1.015743
11	Luteolin	286.24	1.615322
12	6 a hydroxymedicarpin	286.28	0.423520
13	Quercetin	302.24	1.618282
14	Kaempferol 3 sulphate	365.29	1.098278
15	Daidzin	416.38	0.917642
16	Genistin	432.38	1.078467
17	Kaempferol 3 rhamnoside	434.35	0.975201
18	Calycosin 7 O β D glucoside	446.40	2.134107
19	Kaempferol 3 glucoside	448.38	1.489309
20	Kaempferol 7 O β D glucopyranoside	448.38	0.860512
21	Luteolin 7 glucoside	448.38	1.380228
22	Isoquercitrin	464.38	1.559954
23	Hyperoside	464.38	1.535424
24	Querciturone	478.36	0.855362


Based on PASS online server analysis, there are 7 compounds of flavonoid groups' potential to cure the acne (Table 4). Especially the compounds of flavonoid groups contained in water extraction of *D. esculentum*.

Table 4 shows that there are 7 of 13 flavonoid groups potential as antiacne based upon the database of PASS online server indicated with anti-seborrheic, AR expression inhibitor and CYP1A1 inhibitor. The compounds are apigenin, genistein, daidzein, biochanin A, biochanin B, calycosin and quercetin, as antiseborrheic, AR expression inhibitor, and CYP1A1 inhibitor, with PA \geq 0.7. Previous study has shown that apigenin inhibits inflammatory response caused by *P. acnes* [10]. *P. acnes* has lypolitic ability since lipase enzyme can hydrolyze fat to fatty acid and glycerol. Skin tissue of sebaceous gland hyperkeratinizes, blocks the pores, and develops the pustules [30]. Other potential compounds are genistein and daidzein clinically examined to be able to treat inflammation from acne [18, 23, 34]. Quercetin through *in vitro* test also has activity to fight against bacteria *P. acnes* [16].


Table 4. Biological activity of *D. esculentum* flavonoid based on PASS online server analysis

No	Compounds	Probability activity (PA)		
		Anti seborrheic	AR expression inhibitor	CYP1A1 inhibitor
1	Apigenin	0.806	0.773	0.745
2	Genistein	0.832	0.872	0.850
3	Daidzein	0.835	0.831	0.756
4	Naringenin	0.811	-	0.894
5	9 O methylcoumestrol	0.780	0.833	-
6	Biochanin A	0.803	0.869	0.947
7	Biochanin B	0.807	0.828	0.892
8	Calycosin	0.720	0.871	0.930
9	Coumestrol	0.813	0.836	-
10	Hyperoside	-	0.554	0.323
11	Leucocyanidin	0.794	-	-
12	Luteolin	0.873	0.859	-
13	Quercetin	0.835	0.894	0.916


Furthermore, there are 12 potential compounds only as anti-seborrheic, to reduce excessive sebum release of the tissue. Seborrheic is a chronic inflammatory skin disturbance in puberty, from increased skin lipid produced by sebaceous gland development promoting sebum secretion [12]. Those are apigenin, genistein, daidzein, naringenin, 9 O methylcoumestrol, biochanin A, biochanin B, calycosin, coumestrol, leucocyanidin, luteolin and quercetin (Fig. 1).




Apigenin (a)




Genistein (b)




Daidzein (c)




Naringenin (d)




9-O-Methylcoumestrol (e)




Biochanin A (f)



Biochanin B (g)



Calycosin (h)



Coumestrol (i)




Fig. 1 Flavonoid major class structure: apigenin (a), genistein (b), daidzein (c), naringenin (d), 9 O methylcoumestrol (e), biochanin A (f), biochanin B (g), calycosin (h), coumestrol (i), hyperoside (j), leucocyanidin (k), luteolin (l), and quercetin (m) [22].

Table 3 also shows that there are 10 of 13 flavonoid groups' potential as anti-acne in androgen receptor expression inhibitor indicator. Inhibition of androgen receptor can control the appearance of acne [11]. Ten potential compounds as androgen receptor expression inhibitor in *D. esculentum* extract are apigenin, genistein, daidzein, 9 O methylcoumestrol, biochanin A, biochanin B, calycosin, coumestrol, luteolin and quercetin. Furthermore, there are also 8 of 13 flavonoid compounds of *D. esculentum* extract potential as anti-acne in CYP1A1 inhibitor indicator. CYP1A1 expression in human skin is a key marker of aryl hydrocarbon receptor (AHR) activation that causes the presence of blackheads [7]. Those are apigenin, genistein, daidzein, naringenin, biochanin A, biochanin B, calycosin and quercetin.

Conclusion

Diplazium esculentum extract contains various active compounds that are potential to be developed as new drug for acne treatment. This result also reflects that water could be used as more economic and safe alternative solvent. Different number of compounds obtained in both extraction methods has indicated higher effectivity of decoction method than infusa method. In addition, this study could become early step of traditional medicine development using natural material and method.

Acknowledgments

This research was funded by the Indonesian Ministry of Technology and Higher Education Research through the BPP-DN program.

References

1. Alghazeer R., A. Elmansori, M. Sidati, F. Gammoudi, S. Azwai, H. Naas, A. Garbaj, I. Eldaghayes (2017). *In vitro* Antibacterial Activity of Flavonoid Extracts of Two Selected Libyan Algae against Multi-drug Resistant Bacteria Isolated from Food Products, *Journal of Biosciences and Medicines*, 5, 26-48.
2. Azwanida N. N. (2015). A Review on the Extraction Methods Use in Medicinal Plants, Principle, Strength and Limitation, *Medicinal and Aromatic Plants*, 4(3), 1-6.

3. Benmeziane F., R. Djamaï, Y. Cadot, R. Seridi (2014). Optimization of Extraction Parameters of Phenolic Compounds from Algerian Fresh Table Grapes, (*Vitis Vinifera*), Int Food Res J, 21(3), 1061-1065.
4. Chai T. T., M. T. Kwek, H. C. Ong, F. C. Wong (2015). Water Fraction of Edible Medicinal Fern *Stenochlaena palustris* is a Potent α -glucosidase Inhibitor with Concurrent Antioxidant Activity, Food Chemistry, 186, 26-31.
5. Contassot E., L. E. French (2014). New Insights into Acne Pathogenesis: *Propionibacterium acnes* Activates the Inflammasome, Journal of Investigative Dermatology, 132(2), 310-313.
6. Cushnie T. P. T., A. J. Lamb (2005). Antimicrobial Activity of Flavonoids, Int J Antimicrob Agents, 27(2), 181, <http://www.ncbi.nlm.nih.gov/pubmed/16323269>.
7. Fabbrocini G., G. Kaya, S. P. Caseiro, D. Vita, A. Kaya, F. Fontao, O. Sorg, J. H. Saurat (2015). Aryl Hydrocarbon Receptor Activation in *Acne vulgaris* Skin: A Case Series from the Region of Naples, Italy, Dermatology, 231(4), 334-338.
8. Ferreira O., S. P. Pinho (2012). Solubility of Flavonoid in Pure Solvents, Industrial and Engineering Chemistry Research, 51(18), 6586-6590.
9. Fotakis C., D. Tsigirmani, T. Tsiaka, D. Z. Lantzouraki, I. F. Strati, C. Markis, D. Tagkouli, C. Proestos, V. J. Sinanoglou, P. Zoumpoulakis (2016). Metabolic and Antioxidant Profiles of Herbal Infusion and Decoction, Food Chem, 211, 963-971.
10. Huang W. C., T. H. Tsai, C. J. Huang, Y. Y. Li, J. H. Chyuan, L. T. Chuang, P. J. Tsai (2015). Inhibitory Effects of Wild Bitter Melon Leaf Extract on *Propionibacterium acnes*-induced Skin Inflammation in Mice and Cytokine Production *in vitro*, Food Funct, 6(8), 2550-2560.
11. Inoue T., Y. Miki, S. Kakuo, A. Hachiya, T. Kitahara, S. Aiba, C. C. Zouboulis, H. Sasano (2014). Expression of Steroidogenic Enzymes in Human Sebaceous Glands, Bioscientifica, 222(3), 301-312.
12. Del Rosso J. Q. (2011). Adult Seborrheic Dermatitis: A Status Report on Practical Topical Management, Journal of Clinical and Aesthetic Dermatology, 4(5), 32-38.
13. Junejo J. A., G. Gogoi, J. Islam, M. Rudrapal, P. Mondal, H. Hazarika, K. Zaman (2018). Exploration of Antioxidant, Antidiabetic and Hepaprotective Activity of *Diplazium eculentum* – a Wild Edible Plant from North Eastern India, Future Journal of Pharmaceutical Sciences, 4(1), 93-101.
14. Kavak D. D. (2017). Optimization of Extraction Time, Temperature and Solvent Concentration for the Antioxidant Activity and Total Phenolic Content of the *Cydonia oblonga* Mill. Leaves, American-Eurasian Journal of Sustainable Agriculture, 11(6), 1-6.
15. Kim N. H., S. H. Jang, S. H. Kim, H. J. Lee, Y. Kim, J. H. Ryu, M. S. Rhee (2015). Use of Phytic Acid and Hyper-salting to Eliminate *Escherichia coli* O157:H7 from Napa Cabbage for Kimchi Production in a Commercial Plant, International Journal of Food Microbiology, 214(2), 24-30.
16. Kim S., S. Oh, H. B. Noh, et al. (2018). *In vitro* Antioxidant and Anti-*propionibacterium acnes* Activities of Cold Water, Hot Water, and Methanol Extracts, and Their Respective Ethyl Acetate Fractions, from *Sanguisorba officinalis* L. Roots, Molecules, 23(3001), 1-17.
17. Kumar S., A. K. Pandey (2013). Chemistry and Biological Activities of Flavonoids: An Overview, The Scientific World Journal, 1-16, <http://dx.doi.org/10.1155/2013/162750>.
18. Matsumoto T., Y. Matsubara, Y. Mizuhara, et al. (2015). Plasma Pharmacokinetics of Polyphenols in a Traditional Japanese Medicine, Jumihaidokuto, which Suppresses *Propionibacterium acnes*-induced Dermatitis in Rats, Molecules, 20, 18031-18046.
19. Ng'uni T., T. Mothlalamme, R. Daniels, J. Klaasen, B. C. Fielding (2015). Additive Antibacterial Activity of Naringenin Antibiotic Combinations against Multidrug Resistant *Staphylococcus aureus*, Academic Journals, 9(23), 1514-1518.

20. PASS online server, <http://www.pharmaexpert.ru/passonline/> (Access Date June 6, 2022).
21. Paul B., B. Bhuyan, D. D. Purkayastha, S. S. Dhar (2015). Green Synthesis of Silver Nanoparticles Using Dried Biomass of *Diplazium esculentum* (retz.) sw. and Studies of Their Photocatalytic and Anticoagulative Activities, *Journal of Molecular Liquids*, 212, 813-817.
22. Pubchem, <https://pubchem.ncbi.nlm.nih.gov/> (Access Date June 6, 2022).
23. Riyanto P., P. Subchan, R. Lelyana (2015). Advantage of Soybean Isoflavone as Antiandrogen on *Acne vulgaris*, *Dermato Endocrinology*, 7(1), 1-5.
24. Roy S., T. P. Chaudhuri (2017). Toxicological Assessment of *Diplazium esculentum* on the Reproductive Functions of Male Swiss Albino Mouse, *Drug and Chemical Toxicology*, 40(2), 171-182.
25. Septaningsih D. A., L. K. Darusman, F. M. Afendi, R. Heryanto (2018). Liquid Chromatography Mass Spectrometry (LC-MS) Fingerprint Combined with Chemometrics for Identification of Metabolites Content and Biological Activities of *Curcuma aeruginosa*, *Indonesian Journal of Chemistry*, 18(1), 43, <https://doi.org/10.22146/ijc.25456>.
26. Sivasothy Y., S. F. Sulaiman, K. L. Ooi, H. Ibrahim, K. Awang (2013). Antioxidant and Antibacterial Activities of Flavonoids and Curcuminoids from *Zingiber spectabile* Griff, *Food Control*, 30, 714-720.
27. Su Y., L. Ma, Y. Wen, H. Wang, S. Zhang (2014). Studies of the *in vitro* Antibacterial Activities of Several Polyphenols against Clinical Isolates of Methicillin-resistant *Staphylococcus aureus*, *Molecules*, 19, 12630-12639.
28. Swain S. S., R. N. Padhy (2015). *In vitro* Antibacterial Efficacy of Plants Used by an Indian Aboriginal Tribe against Pathogenic Bacteria Isolated from Clinical Samples, *Journal of Taibah University Medical Sciences*, 10(4), 379-390.
29. Tarahovsky Y. S., Y. A. Kim, E. A. Yagolnik, E. N. Muzafarov (2014). Flavonoid-membrane Interactions: Involvement of Flavonoid-metal Complexes in Raft Signalling, *Biochimica et Biophysica Acta*, 1838, 1235-1246.
30. Taylor M., M. M. L. Gonzalez, R. M. Porter (2011). Pathways to Inflammation: Acne Pathophysiology, *European Journal of Dermatology*, 21, 323-333.
31. Wadhwa R., R. Singh, R. Gao, N. Shah, N. Widodo, T. Nakamoto, S. C. Kaul (2013). Water Extract of Ashwagandha Leaves has Anticancer Activity: Identification of an Active Component and Its Mechanism of Action, *PLoS ONE*, 8(10), 1-11.
32. Xiao J., G. Kai (2012). A Review of Dietary Polyphenol-plasma Protein Interactions: Characterization, Influence on the Bioactivity, and Structure-affinity Relationship, *Critical Reviews in Food Science and Nutrition*, 52(1), 85-101.
33. Xie Y., W. Yang, F. Tang, X. Chen, L. Ren (2015). Antibacterial Activities of Flavonoids: Structure Activity Relationship and Mechanism, *Current Medicinal Chemistry*, 22, 132-149.
34. Zaenglein A. L. (2008). *Acne vulgaris* and Acneiform Eruption, *Fitzpatrick's Dermatology in General Medicine*, 7th Ed., New York, McGraw-Hill.
35. Zannah F., M. Amin, H. Suwono, B. Lukiati (2017). Phytochemical Screening of *Diplazium esculentum* as Medicinal Plant from Central Kalimantan, Indonesia, *AIP Conference Proceedings*, Vol. 1844, American Institute of Physics Inc., 1-4.
36. Zannah F., M. Amin, H. Suwono, B. Lukiati (2017). The Utilization of Traditional Medicinal Plants by Dayak Community in Wetland Environment of Central Kalimantan, *Nasional Seminar of Lambung Mangkurat University*, 493-496.

Fathul Zannah, Ph.D.E-mail: fathulzannah@umpalangkaraya.ac.id

Fathul Zannah received her S.Pd. Degree in Biology and M.Pd. Degree from Lambung Mangkurat University, Indonesia, respectively in 2010 and 2012, and Ph.D. Degree from State University of Malang, Indonesia in 2019.

Prof. Mohamad Amin, Ph.D.E-mail: mohamad.amin.fmipa@um.ac.id

Mohamad Amin is a Professor at the College of Biology Education in State University of Malang, Indonesia. He received his S.Pd. Degree in Biology in 1991, M.Sc. Degree in 1997, and Ph.D. Degree in Genetics and Biology Molecular in 2003 from Martin Luther University, Germany.

Prof. Hadi Suwono, Ph.D.E-mail: hadi.suwono.fmipa@um.ac.id

Hadi Suwono is a Lecturer at the College of Biology Education in State University of Malang, Indonesia. He received his S.Pd. Degree of Biology in 1990, M.Sc. Degree in 1993, and Ph.D. Degree in Biology in 2007 from State University of Malang, Indonesia.

Betty Lukiati, Ph.D.E-mail: betty.lukiati.fmipa@um.ac.id

Betty Lukiati is a Lecturer at the College of Biology Education in State University of Malang, Indonesia. She received her S.Pd. Degree of Biology in 1981, M.Sc. Degree in 1990, and Ph.D. Degree in Science in 2012 from Airlangga University, Indonesia.



© 2022 by the authors. Licensee Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).